

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/115847/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Mullish, Benjamin H., McDonald, Julie A.K., Thursz, Mark R. and Marchesi, Julian R. ORCID: <https://orcid.org/0000-0002-7994-5239> 2018. Antibiotic-associated disruption of microbiota composition and function in cirrhosis is restored by fecal transplant. *Hepatology* 68 (3) , p. 1205. 10.1002/hep.30090 file

Publishers page: <http://dx.doi.org/10.1002/hep.30090>
<<http://dx.doi.org/10.1002/hep.30090>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.
See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Correspondence: Antibiotic-Associated Disruption of Microbiota Composition and Function in Cirrhosis is Restored by Fecal Transplant

Benjamin H Mullish¹, b.mullish@imperial.ac.uk

Julie A K McDonald¹, julie.mcdonald@imperial.ac.uk

Mark R Thursz¹, m.thursz@imperial.ac.uk

Julian R Marchesi^{1,2}, j.marchesi@imperial.ac.uk

¹Liver Unit/ Division of Integrative Systems Medicine and Digestive Disease, St Mary's Hospital Campus, Imperial College London

²School of Biosciences, Cardiff University

Keywords: microbiome; metabonome; cirrhosis; antibiotics

To the Editor:

The study from Bajaj and colleagues – evaluating gut microbiome-metabonome changes in cirrhotic patients receiving fecal microbiota transplant (FMT)¹ – merits further discussion of its conclusions.

In this study, cirrhotic patients with hepatic encephalopathy (HE) taking standard of care (SOC) therapy were randomised to either continue with SOC alone, or to also receive five days of antibiotics prior to a single FMT enema. The investigators demonstrate partial recovery of microbiota, bile acid, and short-chain fatty acid (SCFA) profiles for patients in the FMT arm, and conclude that FMT directly restores gut microbiome-metabonome interactions in antibiotic-treated cirrhotics¹. However, there was no arm in this study where patients received antibiotics but no FMT; therefore, it is impossible to distinguish whether any changes observed were directly caused by FMT as claimed, or at least partly represent purely gut microbiota recovery after completion of antibiotics.

There are certainly grounds for questioning the efficacy of the FMT. Specifically, despite selecting a stool donor with high relative abundance of *Lachnospiraceae* and *Ruminococcaceae*, FMT recipients actually demonstrate a significantly reduced relative abundance of these bacterial families post-FMT (day 20) compared to baseline values (day 0) (Figure 2¹). Furthermore, it is well-established that human gut microbiota structure recovers to a profile comparable to that found pre-antibiotics within a short period following completion of an antibiotic course². Additionally, whilst poorly-explored in cirrhosis, SCFA³ and bile acid profiles⁴ reconstitute over a time course of days to weeks after antibiotic cessation in healthy rodents without any specific intervention.

If FMT directly treated HE through modulation of gut microbiota-metabonome interactions, then it would be expected that at least some of these variables would exceed baseline values post-FMT, rather than only match baseline levels at best, as shown. Adverse events associated with FMT are uncommon, but serious adverse events are described – including bacteraemia – in the non-cirrhotic population⁵; such concerns are clearly amplified in decompensated cirrhotics, given this group's propensity to and outcomes from sepsis. Until the safety profile of FMT within cirrhotic patients has been further evaluated – and a potential mechanistic link between FMT and improvement in HE defined - then FMT remains at too early a stage to consider more widely as HE treatment.

References:

1. Bajaj JS, Kakiyama G, Savidge T, et al. Antibiotic-Associated Disruption of Microbiota Composition and Function in Cirrhosis is Restored by Fecal Transplant. *Hepatology*. doi:10.1002/hep.30037.
2. De La Cochetiere MF, Durand T, Lepage P, Bourreille A, Galmiche JP, Dore J. Resilience of the Dominant Human Fecal Microbiota upon Short-Course Antibiotic Challenge. *J Clin Microbiol*. 2005;43(11):5588-5592.
3. **Zhao X, Jiang Z**, Yang F, et al. Sensitive and Simplified Detection of Antibiotic Influence on the Dynamic and Versatile Changes of Fecal Short-Chain Fatty Acids. Virolle M-J, ed. *PLoS One*. 2016;11(12):e0167032.
4. Theriot CM, Koenigsknecht MJ, Carlson PE, et al. Antibiotic-induced shifts in the mouse gut microbiome and metabolome increase susceptibility to *Clostridium difficile* infection. *Nat Commun*. 2014;5:ncomms4114.
5. Baxter M, Colville A. Adverse events in faecal microbiota transplant: a review of the literature. *J Hosp Infect*. 2016;92(2):117-127.

Corresponding author: Dr Benjamin Mullish, Liver Unit/ Division of Integrative Systems Medicine and Digestive Disease, 10th Floor QEQM Wing, St Mary's Hospital Campus, Imperial College London, South Wharf Road, Paddington, London, UK. Tel: +44(0)2033126454; b.mullish@imperial.ac.uk .

List of abbreviations:

FMT fecal microbiota transplant

HE hepatic encephalopathy

SOC standard of care

SCFA short chain fatty acids

Financial support: The Division receives financial support from the National Institute of Health Research (NIHR) Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial College London. BHM is the recipient of a Medical Research Council Clinical Research Training Fellowship (grant reference: MR/R000875/1).